

Riverine Carbon and the Sedimentary Record on the Continental Shelves

Stefano Miserocchi and Leonardo Langone
Istituto Scienze Marine, Sezione Geologia Marina
(formerly Istituto di Geologia Marina)

Consiglio Nazionale delle Ricerche

Via Gobetti, 101

40129 Bologna, Italy

phone: +39 (051) 6398880 Fax. +39 (051) 6398940 email: stefano.miserocchi@ismar.cnr.it

Award Number: N00014-03-1-0154

LONG-TERM GOALS

The aim of our contribution (as part of the EuroSTRATAFORM Program) was to improve a more complete understanding of organic matter (OM) transport, deposition and burial in the Gulf of Lions and Adriatic Sea. Continental shelves represent the largest modern repository of particulate organic carbon (OC); nevertheless, some major river systems may be sites where a substantial portion of terrestrial inputs bypass the shelf, and are discharged onto the continental slope and beyond. Characterizing the origin of OM delivered to the ocean and defining the processes that affect its ultimate burial may provide a better understanding of the mechanisms that influence organic carbon preservation in the ocean.

OBJECTIVES

During this year our research was mainly focused on the Gulf of Lions: collecting sediment samples and carrying out elemental and stable carbon isotopic analysis. The field work was carried out in the Gulf of Lions taking into account three different depositional elements: the prodelta Rhône area, the continental shelf, and the array of submarine canyons, which incise the shelf and the slope. Our approach was to use the signature of the organic matter in order to:

- 1) trace the initial distribution of terrigenous material accumulated in the prodelta Rhône area and follow its transport along the continental shelf;
- 2) characterize qualitatively the riverine input defining the amount of soil and woody debris, respectively;
- 3) define the amount of riverine OC sequestered by the canyons and exported towards abyssal plain;
- 4) verify the influence of minor rivers besides the Rhône river (such as the Têt river).

APPROACH

Terrestrial and marine organic matter burial in marine sediments is an important reservoir, representing the predominant long-term sink in the global biogeochemical cycle of OC (Berner, 1989). Deltas and

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Riverine Carbon and the Sedimentary Record on the Continental Shelves			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Istituto Scienze Marine, Sezione Geologia Marina,(formerly Istituto di Geologia Marina),Consiglio Nazionale delle Ricerche,Via Gobetti, 101,40129 Bologna, Italy, ,			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The aim of our contribution (as part of the EuroSTRATAFORM Program) was to improve a more complete understanding of organic matter (OM) transport, deposition and burial in the Gulf of Lions and Adriatic Sea. Continental shelves represent the largest modern repository of particulate organic carbon (OC); nevertheless, some major river systems may be sites where a substantial portion of terrestrial inputs bypass the shelf, and are discharged onto the continental slope and beyond. Characterizing the origin of OM delivered to the ocean and defining the processes that affect its ultimate burial may provide a better understanding of the mechanisms that influence organic carbon preservation in the ocean.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

continental shelves, considered to be the main repositories for organic carbon in marine sediments, receive large inputs from both autochthonous and allochthonous sources and account for ~90% of the OC burial in the ocean (Hedges and Keil, 1995). However, these same massive storage systems are often sites of efficient remineralization (Aller and Blair, 2004): some studies suggest that in highly reactive mobile shelf muds, physical reworking, bioturbation, exposure, and reoxidation inhibit high carbon burial (Aller, 1998, Aller and Blair, 2004). In the Gulf of Lions and the Adriatic Sea a simple two-end-member mixing model, in which vascular plant debris is diluted by marine phytoplankton detritus, can describe the composition of OM in coastal marine sediments. The isotopic signature and elemental composition of the sedimentary OC are widely used as proxies for decoding origin and fate of OM on various continental margins. Vascular plants of terrestrial origin exhibit many bulk chemical and isotopic proprieties that distinguish them from marine organisms. For example, the predominance of nitrogen-free biomacromolecules (e.g. tannin, lignin, cellulose, cutin and suberin) over proteins (C:N \approx 3-4) makes higher plant tissues characteristically carbon-rich (C:N \approx 20-500) versus phytoplankton (C:N \approx 6.6) and bacteria (C:N \approx 4). The organic matter absorbed by soil shows a value between vascular plants and bacteria (C:N \approx 8-14). Carbon isotopic signatures ($\delta^{13}\text{C}$) can distinguish organic matter photosynthesized by C3 plants (from -25 up to -28 ‰) from temperate marine phytoplankton (from -18‰ up to -22‰) (Hedges et al., 1997).

WORK COMPLETED

Concerning the Gulf of Lions, box cores were collected:

- in the Têt and Rhône delta along shore-normal transects;
- near the heads of Cap de Creus and Lacaze-Duthiers Canyons along transects, as well as a smaller number in the deeper portion of these canyons;
- along a mud wedge on the shelf following the bathymetry of 75 m;

Vertical sediment profiles from selected box and kasten cores were also collected. The samples were collected in October 2004 on *R/V Oceanus* and in February and May 2005 on *R/V Endeavor*. The uppermost and the lowest sediment slices were subsampled from each box-core.

Concerning the Adriatic Sea, we completed the OC analyses. For both the Adriatic Sea and the Gulf of Lions, approximately 750 samples were collected and analyzed for carbon stable-isotopes, OC, and total N determination at ISMAR, using a FINNIGAN Delta Plus mass spectrometer directly coupled to the FISOONS NA2000 Element Analyzer.

RESULTS

Gulf of Lions.

Fig. 1 shows the OC content on surface sediments in the GOL. The OC exhibits the highest values in the prodelta Rhône area up to 1.44 %. On going offshore, the OC content decreases rapidly down to 0.46 %. In contrast, the Têt prodelta exhibits lower values ranging from 0.29 to 0.88 %. Consequently, the Rhône plays a much more important role than the Têt in the flux of OC to the sea floor. The sedimentary OC in the prodelta Rhône area is isotopically depleted compared to the material

accumulated on the rest of the shelf (Fig. 2). These depleted values in $\delta^{13}\text{C}$ indicate a clear terrestrial OM origin that moving offshore is diluted by marine phytoplankton detritus. The “refractory” organic material delivered by the Rhône, being partially degraded, especially in the soil fraction, is more efficiently accumulated and preserved in the surficial sediment than fresh and “labile” algal compounds. For that reason, the OC content exhibits a drastic drop-off when going offshore far away from any direct terrestrial input. The high correlation ($r^2 = 0.91$) among OC content and $\delta^{13}\text{C}$ along a transect across-shelf confirms this (Fig. 3).

The compositional characteristics of OM in the Cap de Creus Canyon indicate a primarily marine source. The samples on the shelf close to the canyon head are still slightly terrestrial, showing isotopically depleted $\delta^{13}\text{C}$ values, whereas the deepest samples in the canyon are plainly marine. Based on these OC contour maps, it seems that the organic material from the Rhône prodelta is driven westwards by the LPC (Liguro-Provençal Current), and on its way the OC is degraded efficiently, showing low concentrations (up to 0.10%) at the head of the Cap de Creus canyon.




Figure 1. Distribution of OC content (% wt) on surface sediment in the shelf of Gulf of Lions and in the Cap de Creus Canyon. White crosshairs indicate station locations.




Figure 2. Distribution of $\delta^{13}\text{OC}$ (‰) in surface sediment on the shelf of Gulf of Lions and in the Cap de Creus Canyon. White crosshairs indicate station locations.




Figure 3. OC content (% OC) and its isotopic OC composition $\delta^{13}\text{OC}$ ‰) in surface sediment along a transect across-shelf from the prodelta Rhône area up to shelf break. [The strong correlation ($r^2=0.91$) indicates the highest contents are associated to the terrigenous material].

Adriatic Sea.

Fig. 4 shows the OC content in surface sediments on the western Adriatic continental shelf. In the Po prodelta, the high primary productivity and the considerable amount of OC delivered by the Po River directly affect the surface sediments. Organic carbon content higher than 1% composes a crushed depocenter developing around the delta, probably as a result of high and rapid sediment accumulation enhanced by large rapidly sinking flocs (Fox et al., 2004), which prevent grain-size sorting. Furthermore, four smaller isolated depocenters, just in front of each distributary, seem to underline the key role of river load on surface marine sediments.

On going southward, the region closest to the Po seems to remain under the influence of organic material supplied to the Po River, which spreads southwards driven by the anticyclonic circulation.

Beyond this region, on the shelf, OC values increase offshore. Because of water dynamics, the river sediment load delivered to the sea is hydraulically sorted in grain size, in accordance with the classic modern sedimentation pattern on continental shelves: costal sands, mid-shelf muds and relict shelf sands further offshore (Brambati et al., 1983). The deposition of fine sediment, which generally has elevated levels of associated OM relative to their coarse counterparts (Hedges and Keil, 1995), strongly influences the OC distribution in surface sediments.




Figure 4. *Distribution of OC content (% wt) on surface sediment in the western Adriatic Continental shelf. Black crosshairs indicate station locations.*

IMPACT/APPLICATIONS

Studying the sedimentary processes, which affect the distribution and preservation of organic matter on the continental shelves will provide key insights to understanding the role this environment as sinks for OC with respect to the global carbon cycle.

RELATED PROJECTS

Our research is related to other groups. First we are collaborating with T. Milligan (BIO), in order to examine the relationship between organic carbon and particle grain size.

Second, we are working with D. Orange and A. Garcia (UCSC) in evaluating the origin of anomalous sub-surface shallow gas concentrations in the Rhône prodelta delta and in the Cap de Creus Canyon. Our hypothesis is that, away from the flood-related sediments off the Rhône delta (where methane content is up to 87440 ppm), the organic matter is being reworked and remineralized on its way down the western coast of the Gulf of Lions, so that the recent deposits in the canyon contain little reactive carbon.

Finally, we are cooperating with J. Fabres and A. Calafat (CRG Marine Geosciences) in carrying out elemental and isotopic analysis on suspended material collected with sediment traps deployed in all canyons of GOL. The suspended material collected during flood events (Rhône and Têt) showed a

clear terrestrial source, improving the understanding of the factors that control sediment transfer across shelf breaks.

REFERENCES

Aller, R.C, 1998. Mobile deltaic and continental shelf muds as suboxic, fluidized bed reactors. *Marine Chemistry* 61, 143-155.

Aller, R.C, Blair, N.E., 2004. Early diagenetic remineralization of sedimentary organic C in the Gulf of Papua deltaic complex (Papua New Guinea): Net loss of terrestrial C and diagenetic fractionation of C isotopes. *Geochimica et Cosmochimica Acta* 68, 1815-1825.

Berner, R.A., 1989. Biogeochemical cycles of carbon and sulfur and their effect on atmospheric oxygen over phanerozoic time. *Palaeogeography, Palaeoclimatology, Palaeoecology* 75, 97–122.

Brambati, A., Ciabatti, M., Fanzutti, G.P., Marabini, F., Marocco, R., 1983. A new sedimentological textural map of the northern and central Adriatic Sea. *Bollettino di Oceanografia Teorica e Applicata* I, 267-271.

Hedges, J.I., Keil, R.G., 1995. Sedimentary organic matter preservation: an assessment and speculative synthesis. *Marine Chemistry* 49, 81–115

Hedges, J.I., Keil, R.G., Benner, R., 1997. What happens to terrestrial organic matter in the ocean? *Organic Geochemistry* 27, 195–212.

Fox, J.M., Hill, P.S., Milligan T.G., Boldrin, A., 2004. Flocculation and sedimentation on the Po River Delta. *Marine Geology* 203, 95-107.

PUBLICATIONS

Papers

Boldrin, A., Langone, L., Miserocchi S., Turchetto M., Acri A. Po River plume on the Adriatic continental shelf: dispersion and sedimentation of dissolved and suspended matter during different river discharge rates. *Marine Geology* [in press, refereed].

Cattaneo A., Trincardi F., Langone L., Asioli A., and Puig P., 2004. Clinoform generation on Mediterranean margins. *Oceanography* 17, 104-117.

Fabres J., Calafat A., Tesi T., Zúñiga D., Canals M., Heussner S., Miserocchi S. , X. Durrieu de Madron, Bourrin F., Palanques A., Puig P. The impact of a winter catastrophic storm on near bottom particle fluxes at canyon heads in the Gulf of Lions. *Marine Geology* [Submitted].

Garcia-Garcia A., Orange D., Miserocchi S., Correggiari A., Langone L., Lorenson T., Trincardi F., Nittrouer C.A. What controls shallow gas in Western Adriatic Sea? *Continental Shelf Research*. [Submitted].

Garcia-Garcia A., Orange D., Lorenson, T., Radakovitch, O., Tesi, T., Miserocchi, S., Berne, S., Friend, P. L., Nittrouer, C., Normand, A. Shallow gas studies off the Rhône prodelta, Gulf of Lions. Marine Geology [Submitted].

García-García, A., Orange, D., Lorenson, T., Tesi, T., Herbert, I., Miserocchi, S., Sansoucy, M., Dougherty, J. Cap de Creus canyon: a canyon with active sediment accumulation and little shallow gas. Marine Geology. [Submitted].

Miserocchi S., Langone L., Tesi T. Concentration and isotopic composition of organic carbon within a flood layer in the Po River prodelta (Adriatic Sea). Continental Shelf Research [Submitted].

Nittrouer C.A., Miserocchi S., Tricardi F., 2004. The PASTA Project: investigation of Po and Apennine Sediment Transport and Accumulation. Oceanography 17, 46-57.

Orange D., García-García A., Lorenson T., Nittrouer C., Milligan T., Miserocchi S., Langone L., Correggiari A., and Trincardi F. Shallow gas and flood deposition on the Po Delta. Marine Geology [in press, refereed].

Tesi T., Miserocchi S., Langone L., Boni L., Guerrini F. Sources, fate and distribution of organic matter on the western Adriatic continental shelf, Italy. Water, Air & Soil Pollution. [Submitted]

Abstract

Fain A., Ogston A.S., Miserocchi S., Palinkas C.M. 2004. Sediment dispersal mechanisms on the Western Adriatic continental shelf. AGU-Ocean Sciences meeting 26-30 January 2004 Portland, Oregon, USA Eos Trans. AGU, 84., (52), Ocean Sci. Meet. Suppl., Abstract OS51B-01, 2003

Miserocchi S., Langone L., Palinkas C., Nittrouer C.A. (2003). Dispersal of riverine organic carbon along the western Adriatic continental shelf. ComDelta -Comparing Mediterranean and Black Sea Prodeltas, 26-28 October 2003, Aix-en-Provence, France

Miserocchi S., Langone L., Palinkas C., Nittrouer C. A. (2003). Organic Carbon and its Isotopic Composition in Surficial Sediments from the Western Adriatic Continental Shelf. AGU Fall Meeting, San Francisco, USA, 8-12 Dec 2003.

Miserocchi S., Langone L., Boldrin A., Turchetto M., Tesi T. 2004. Dispersione del carbonio organico di origine fluviale sul lato occidentale della piattaforma continentale adriatica. Ciclo del Carbonio e Risorse del mare Workshop dell'Istituto di Scienze Marine ISMAR, CNR, Roma, 10-11 maggio 2004.

Miserocchi S., Langone L., Tesi T. (2004) Characterizing sedimentary organic carbon within a flood layer in the Po River prodelta. 32nd International Geological Congress, Florence, Italy.

Miserocchi S., Langone L., Tesi T. (2005) Characterizing sedimentary organic carbon within a flood layer in the Po River prodelta. EuroStrataform meeting-2005. Venice, Italy.

Miserocchi S., Langone L., Tesi T. (2005). Source and distribution of organic matter on the western Adriatic Continental Shelf. 10th International Symposium on the Interactions Between Sediments and Water - IASWS2005. Bled, Slovenia.

Tesi T., Langone L., Miserocchi S. (2005). Organic carbon and its isotopic composition in surficial sediments on the western Adriatic Continental, Italy. Summer Aslo meeting 2005, Santiago de Compostela. Spain.